

## **TECHNICAL DATA SHEET**

## CHYMOSIN LIQUID 1000IMCU/ML

**Introduction:** Our Chymosin (Fermentation Produced Chymosin) is a milk clotting enzyme produced by fermentation of *Kluyveromyces lactis* strain. It is used to replace the pure chymosin coming from a milk sucking calf in its first days of life for coagulating milk proteins during cheese-making. Its performance and properties are identical to calf Chymosin.

Specification: Meet the GB/T 1886.174 -2016 (enzymes for food industry)

| Appearance        | Light yellow to light brown liquid | pН                     | 5.0 - 6.0        |
|-------------------|------------------------------------|------------------------|------------------|
| Clotting activity | ≥1000IMCU/ml                       | Total plate count      | ≤1,000 CFU/ml    |
| Odor & Taste      | Slight odor of fermentation        | Coliform bacteria      | <30 CFU/ml       |
| Specific gravity  | 1.08-1.12g/cm <sup>3</sup>         | Mold & Yeast count     | ≤10CFU/ml        |
| Lead (Pb)         | ≤3 ppm                             | Staphylococcus aureus  | Negative in 1ml  |
| Arsenic (As)      | ≤1 ppm                             | Salmonella spp.        | Negative in 25ml |
| Mercury (Hg)      | ≤0.5 ppm                           | Listeria monocytogenes | Negative in 25ml |
| Cadmium (Cd)      | ≤0.5 ppm                           | E. coli                | Negative in 25ml |

**Characteristics:** Chymosin is a little bit salty, good stability, sodium benzoate free, reliable coagulation, and suitable for the production of various types of cheese with high yields and cheese quality.

Composition: water, glycerin, Chymosin from Kluyveromyces lactis, sodium chloride, trehalose.

**Dosage**: 45 to 50 ml is recommended for every 1,000 liters of milk. The product should be dissolved in pasteurized, chilled and chlorine-free water. Mix with the milk, stirring for 5 minutes.

Packaging: 20L/barrel, 25L/barrel.

**Shelf life and storage**: 1 year when stored in a cool, dry place between 4-8°C, away from sunlight. The product used after the above date has not harmful effects but only a potential decreased activity. Store the original sealed containers at 4-8°C, after the arrival in your factory.